

Obstetrics & Gynecology Science

Yutaka Yoneoka, et al. Recurrence risk of patients with cervical cancer

Supplementary Table 1. Characteristics of low-risk group

Characteristics	L-SCC group (n=38)	L-Adeno group (n=38)	P-value
Age (yr)	51 (38–68)	53 (42–82)	0.43
BMI (kg/m ²)	20.0 (17.6–30.9)	19.7 (16.7–28.7)	0.57
Tumor size (cm)			0.14
≤4	38 (100.0)	37 (97.4)	
>4	0 (0.0)	1 (2.6)	
Stromal invasion			0.37
≤1/2	30 (78.9)	31 (81.6)	
>1/2	8 (21.1)	7 (18.4)	
LVSI			0.31
Negative	27 (71.1)	29 (76.3)	
Positive	11 (28.9)	9 (23.7)	
Vaginal invasion			0.04
Negative	35 (92.1)	38 (100.0)	
Positive	3 (7.9)	0 (0.0)	

Values are presented as median (range) or number of patients (%).

L-SCC group, low-risk group with squamous cell carcinoma; L-Adeno group, low-risk group with endocervical adenocarcinoma, usual type, or adenosquamous carcinoma; BMI, body mass index; LVSI, lymphovascular space invasion.